

$B_{s(d)}^0 - \bar{B}_{s(d)}^0$ mixing and new physics effects in a top quark two-Higgs doublet model

Lin-xia Lü ^{a,b*} and Zhen-jun Xiao ^{a†}

*a. Department of Physics and Institute of Theoretical Physics,
Nanjing Normal University, Nanjing, Jiangsu 210097, P.R.China and*

*b. Department of Physics, Nanyang Teacher's College,
Nanyang, Henan 473061, P.R.China*

(Dated: February 2, 2008)

Abstract

We calculate the new physics contributions to the neutral B_d^0 and B_s^0 meson mass splitting ΔM_d and ΔM_s induced by the box diagrams involving the charged-Higgs bosons in the top quark two-Higgs doublet model (T2HDM). Using the precision data, we obtain the bounds on the parameter space of the T2HDM: (a) for fixed $M_H = 400$ GeV and $\delta = [0^\circ, 60^\circ]$, the upper bound on $\tan \beta$ is $\tan \beta \leq 30$ after the inclusion of major theoretical uncertainties; (b) for the case of $\tan \beta \leq 20$, a light charged Higgs boson with a mass around 300 GeV is allowed; and (c) the bounds on $\tan \beta$ and M_H are strongly correlated: a smaller (larger) $\tan \beta$ means a lighter (heavier) charged Higgs boson.

PACS numbers: 12.15.Ff, 12.60.Fr, 14.40.Nd, 14.65.Fy

*Electronic address: lulinxia@email.njnu.edu.cn

†Electronic address: xiaozhenjun@njnu.edu.cn

I. INTRODUCTION

As a flavor changing neutral current process, $B_q^0 - \bar{B}_q^0$ mixing with $q = d, s$ are generated at the loop-level and have been of fundamental importance in probing virtual effects from potential new physics beyond the standard model (SM). The $B_q^0 - \bar{B}_q^0$ mixing is responsible for the small mass differences between the heavy and light mass eigenstates of neutral B mesons:

$$\Delta M_q = M_{B_{q,H}^0} - M_{B_{q,L}^0}. \quad (1)$$

The mass splitting ΔM_d has been measured with high precision [1, 2], while the measurement of ΔM_s is very difficult due to the rapid oscillation of B_s meson and has been reported by CDF and D0 Collaboration [3, 4, 5] very recently. The world average for ΔM_d [2] and the first observation of ΔM_s from CDF [4] are the following

$$\Delta M_d = 0.507 \pm 0.005 \text{ ps}^{-1}, \quad (2)$$

$$\Delta M_s = 17.77 \pm 0.10(\text{stat}) \pm 0.07(\text{syst}) \text{ ps}^{-1}, \quad (3)$$

which agree well with the standard model (SM) predictions or the results from global fit [6]. The perfect agreement between the SM prediction and the experimental measurements permit us to put strong constraints on the parameter spaces of various new physics models.

In the SM, $B^0 - \bar{B}^0$ mixing is dominated by the box diagrams with two internal t -quarks and W gauge bosons. In new physics models, the box diagrams with one or two W gauge bosons replaced by the new charged scalars or vector bosons and/or top quarks replaced by new fermions can also contribute to $B^0 - \bar{B}^0$ mixing. Using the precision data, we study both $B_d^0 - \bar{B}_d^0$ and $B_s^0 - \bar{B}_s^0$ mixing in the top quark two-Higgs doublet model (T2HDM) and try to find the constraints on the parameter space of this model.

During the past years, $B^0 - \bar{B}^0$ mixing has been studied extensively in the SM and various new physics models. The charged-Higgs boson contributions to $B^0 - \bar{B}^0$ mixing have been calculated at the leading order (LO) for a long time [7]. The next-to-leading order (NLO) quantum chromodynamics (QCD) correction to $B^0 - \bar{B}^0$ mixing is firstly presented in Ref. [8] and the analytic formulae for the QCD renormalization group factors are given in Ref. [9]. In Ref. [10], the authors studied the new physics effects in the conventional two-Higgs-doublet model (2HDM) of type I and II. In Refs. [11, 12, 13] the authors have calculated the charged Higgs boson contributions to the mass splitting ΔM_{B_d} and drawn the constraints on the parameters of the third type of 2HDM (the model III) at the LO or NLO level. Very recently, $B_s^0 - \bar{B}_s^0$ mixing has been used to put constraints on various new physics models, for example, in Refs. [14] after the release of the new data of ΔM_s . In this paper, we will calculate the new physics contributions to ΔM_q within the framework of T2HDM [15, 16, 17]. By comparing the theoretical predictions with the precision data, we draw the constraints on the free parameters of T2HDM.

The organization of this paper is as follows. In sec. II, we firstly give a brief review for the top quark two-Higgs doublet model, and then present the one-loop contributions to the mass splitting ΔM_d and ΔM_s induced by the box diagrams involving the charged Higgs bosons in the T2HDM. Numerical results are presented in sec. III, and the conclusions are included in the final section.

II. $B^0 - \bar{B}^0$ MIXING IN THE T2HDM

The new physics model considered here is the T2HDM proposed in Ref. [15] and studied for example in Refs. [16, 17, 18], which is also a special case of the 2HDM of type III [19]. This model is designed to accommodate the heaviness of the top quark by coupling it to a scalar doublet with large vacuum expectation value (VEV). All the other five quarks are coupled to another scalar doublet, whose VEV is much smaller. As a result, $\tan \beta$ is naturally large in this model.

Let us now briefly recapitulate some important features of the T2HDM [15]. Consider the Yukawa Lagrangian of the form:

$$\mathcal{L}_Y = -\bar{L}_L \phi_1 E l_R - \bar{Q}_L \phi_1 F d_R - \bar{Q}_L \tilde{\phi}_1 G \mathbf{1}^{(1)} u_R - \bar{Q}_L \tilde{\phi}_2 G \mathbf{1}^{(2)} u_R + H.c. \quad (4)$$

where Q_L and L_L are 3-vector of the left-handed quark and lepton doublets; ϕ_i ($i = 1, 2$) are the two Higgs doublets with $\tilde{\phi}_i = i\tau_2 \phi_i^*$; and E , F and G are the 3×3 matrices in the generation space and give masses respectively to the charged leptons, the down and up type quarks; $\mathbf{1}^{(1)} \equiv \text{diag}(1, 1, 0)$ and $\mathbf{1}^{(2)} \equiv \text{diag}(0, 0, 1)$ are the two orthogonal projection operators onto the first two and the third families respectively.

The Yukawa couplings involving the charged-Higgs bosons are of the form [15]

$$\begin{aligned} \mathcal{L}_Y^C = & \frac{g}{\sqrt{2}M_W} \{ -\bar{u}_L V M_D d_R [G^+ - \tan \beta H^+] + \bar{u}_R M_U V d_L [G^+ - \tan \beta H^+] \\ & + \bar{u}_R \Sigma^\dagger V d_L [\tan \beta + \cot \beta] H^+ + h.c. \}. \end{aligned} \quad (5)$$

where $G^\pm$ and $H^\pm$ denote the would-be Goldstone bosons and the physical charged Higgs bosons, respectively. Here M_U and M_D are the diagonal up- and down-type mass matrices, V is the usual CKM matrix and $\Sigma \equiv M_U U_R^\dagger \mathbf{1}^{(2)} U_R$. $U_R^\dagger$ is the unitary matrix which diagonalizes the right-handed up-type quarks and has the following form:

$$U_R = \begin{pmatrix} \cos \phi & -\sin \phi & 0 \\ \sin \phi & \cos \phi & 0 \\ 0 & 0 & 1 \end{pmatrix} \times \begin{pmatrix} 1 & 0 & 0 \\ 0 & \sqrt{1 - |\epsilon_{ct} \xi|^2} & -\epsilon_{ct} \xi^* \\ 0 & \epsilon_{ct} \xi & \sqrt{1 - |\epsilon_{ct} \xi|^2} \end{pmatrix}. \quad (6)$$

where $\epsilon_{ct} \equiv m_c/m_t$, and $\xi = |\xi|e^{i\delta}$ is a complex number of order unity. Inserting Eq. (6) into the definition of Σ yields

$$\Sigma = \begin{pmatrix} 0 & 0 & 0 \\ 0 & m_c \epsilon_{ct}^2 |\xi|^2 & m_c \epsilon_{ct} \xi^* \sqrt{1 - |\epsilon_{ct} \xi|^2} \\ 0 & m_c \xi \sqrt{1 - |\epsilon_{ct} \xi|^2} & m_t (1 - |\epsilon_{ct} \xi|^2) \end{pmatrix}. \quad (7)$$

Now we are ready to calculate the charged Higgs and would-be Goldstone bosons contributions to the mass splitting $\Delta M_{B_q}(q = d, s)$ in the T2HDM.

The effective weak Hamiltonian for $\Delta B = 2$ processes beyond the SM can be written as

$$\mathcal{H}_{\text{eff}}^{\Delta B=2} = \sum_{i=1}^5 C_i Q_i + \sum_{i=1}^3 \tilde{C}_i \tilde{Q}_i \quad (8)$$

with

$$\begin{aligned}
Q_1 &= (\bar{q}_L^\alpha \gamma^\mu b_L^\alpha) (\bar{q}_L^\beta \gamma_\mu b_L^\beta), \\
Q_2 &= (\bar{q}_R^\alpha b_L^\alpha) (\bar{q}_R^\beta b_L^\beta), \quad Q_3 = (\bar{q}_R^\alpha b_L^\beta) (\bar{q}_R^\beta b_L^\alpha), \\
Q_4 &= (\bar{q}_R^\alpha b_L^\alpha) (\bar{q}_L^\beta b_R^\beta), \quad Q_5 = (\bar{q}_R^\alpha b_L^\beta) (\bar{q}_L^\beta b_R^\alpha),
\end{aligned} \tag{9}$$

where $q = s, d$, corresponding to the operators of B_s and B_d system respectively, $P_{L,R} \equiv (1 \mp \gamma_5)/2$, and α, β are color indexes. The tilde operators $\tilde{Q}_i$ ($i = 1, 2, 3$) correspond to the ones Q_i ($i = 1, 2, 3$) with opposite chirality.

In T2HDM, there are two CP-even scalars (H^0, h^0), one CP-odd scalar (A^0), two charged Higgs bosons ($H^\pm$), and the Goldstone bosons ($G^\pm, G^0$). At one loop level only the charged scalars are relevant for the box diagrams contributing to the $B^0 - \bar{B}^0$ mixing amplitude. From Eq. (5), we can rewrite the vertex couplings of $H_l^+ \equiv (H^+, G^+)$ [20] (where $G^\pm$ is the would-be Goldstone boson) in a compact form

$$\mathcal{L}_{int} = H_l^+ \bar{u}_A V_{AI} (a_L^{AI} P_L + a_R^{AI} P_R) d_I + h.c. \tag{10}$$

where

$$a_L^{AI} = \frac{e}{\sqrt{2}s_W} \frac{m_A}{M_W} \cdot \begin{cases} \left(\frac{(\Sigma^\dagger V)_{AI}}{m_A V_{AI}} - 1 \right) \tan \beta, & \text{for } l=1, \\ 1, & \text{for } l=2, \end{cases} \tag{11}$$

$$a_R^{AI} = \frac{e}{\sqrt{2}s_W} \frac{m_I}{M_W} \cdot \begin{cases} \tan \beta, & \text{for } l=1, \\ -1, & \text{for } l=2, \end{cases} \tag{12}$$

with $m_A = (m_u, m_c, m_t)$ and $m_I = (m_d, m_s, m_b)$. The contributions of H_l^+ to the Wilson coefficients C_i of the relevant operators responsible for $B^0 - \bar{B}^0$ mixing can be easily expressed in terms of the coefficients a_L^{AI} and a_R^{AI} as follows.

The contributions to the Wilson $C_1(\mu)$ and $C_2(\mu)$ induced by the box diagrams with one $W^\pm$ and one $H^\pm$ propagator can be written as ¹:

$$\begin{aligned}
C_1(\mu) &= \frac{e^2}{2s_W^2} \sum_{A,A'} \frac{V_{AI}^* V_{AJ} V_{A'I}^* V_{A'J}}{16\pi^2} m_A m_{A'} a_L^{\dagger AI1} a_L^{A'J1} \cdot D_0(m_A^2, m_{A'}^2, M_W^2, M_{H^+}^2), \\
C_2(\mu) &= \frac{e^2}{2s_W^2} \sum_{A,A'} \sum_{l=1}^2 \frac{V_{AI}^* V_{AJ} V_{A'I}^* V_{A'J}}{16\pi^2} a_R^{\dagger AI l} a_R^{A'J l} \cdot 4D_{00}(m_A^2, m_{A'}^2, M_W^2, m_{H_l^+}^2),
\end{aligned} \tag{13}$$

where $s_W \equiv \sin \theta_W$ (θ_W is the Weinberg angle), while the four-point integral functions D_0

¹ The contribution of $G^\pm$ to $C_1(\mu)$ is already taken into account in the Inami-Lim function $S_0(x_t)$ [21]. Mass of the u quark are neglected.

TABLE I: The "magic numbers" appearing in the calculation of the Wilson coefficients in the process of $B^0 - \bar{B}^0$ mixing.

	1	2	3	4	5		1	2	3	4	5
a_i	0.286	-0.692	0.787	-1.143	0.143						
$b_i^{(11)}$	0.865	0	0	0	0	$c_i^{(11)}$	-0.017	0	0	0	0
$b_i^{(22)}$	0	1.879	0.012	0	0	$c_i^{(22)}$	0	-0.18	-0.003	0	0
$b_i^{(23)}$	0	-0.493	0.18	0	0	$c_i^{(23)}$	0	-0.014	0.008	0	0
$b_i^{(32)}$	0	-0.044	0.035	0	0	$c_i^{(32)}$	0	0.005	-0.012	0	0
$b_i^{(33)}$	0	0.011	0.54	0	0	$c_i^{(33)}$	0	0.000	0.028	0	0
$b_i^{(44)}$	0	0	0	2.87	0	$c_i^{(44)}$	0	0	0	-0.48	0.005
$b_i^{(45)}$	0	0	0	0.961	-0.22	$c_i^{(45)}$	0	0	0	-0.25	-0.006
$b_i^{(54)}$	0	0	0	0.09	0	$c_i^{(54)}$	0	0	0	-0.013	-0.016
$b_i^{(55)}$	0	0	0	0.029	0.863	$c_i^{(55)}$	0	0	0	-0.007	0.019

and D_{00} can be written as

$$\begin{aligned}
D_0(a, b, c, d) &= \int \frac{d^4 q}{i\pi^2} \frac{1}{(q^2 - a)(q^2 - b)(q^2 - c)(q^2 - d)} \\
&= \frac{a}{(b-a)(c-a)(d-a)} \log\left[\frac{a}{d}\right] + \frac{b}{(a-b)(b-c)(b-d)} \log\left[\frac{b}{d}\right] \\
&\quad + \frac{c}{(a-c)(b-c)(d-c)} \log\left[\frac{c}{d}\right], \tag{14}
\end{aligned}$$

$$\begin{aligned}
D_{00}(a, b, c, d) &= \frac{1}{4} \int \frac{d^4 q}{i\pi^2} \frac{q^2}{(q^2 - a)(q^2 - b)(q^2 - c)(q^2 - d)} \\
&= \frac{a^2}{(b-a)(c-a)(d-a)} \log\left[\frac{a}{d}\right] + \frac{b^2}{(a-b)(b-c)(b-d)} \log\left[\frac{b}{d}\right] \\
&\quad + \frac{c^2}{(a-c)(b-c)(d-c)} \log\left[\frac{c}{d}\right]. \tag{15}
\end{aligned}$$

The contributions induced by the box diagrams with two $H_l^\pm$ propagators can be written as ²:

$$\begin{aligned}
C_1(\mu) &= -\frac{1}{2} \sum_{A, A'} \sum_{l, n} \frac{V_{AI}^* V_{AJ} V_{A'I}^* V_{A'J}}{16\pi^2} a_L^{\dagger AI} a_L^{AJn} a_L^{\dagger A'In} a_L^{A'Jl} \\
&\quad \cdot D_{00} \left(m_A^2, m_{A'}^2, M_{H_l^+}^2, M_{H_n^+}^2 \right), \tag{16}
\end{aligned}$$

² In the sum over l and n in the expression for $C_1(\mu)$ the contribution of $G^\pm G^\mp$ is excluded since it has been taken into account in the function $S_0(x_t)$ [21].

$$\tilde{C}_1(\mu) = -\frac{1}{2} \sum_{A,A'} \sum_{l,n}^2 \frac{V_{AI}^* V_{AJ} V_{A'I}^* V_{A'J}}{16\pi^2} a_R^{\dagger AI l} a_R^{AJ n} a_R^{\dagger A' I n} a_R^{A' J l} \cdot D_{00} \left(m_A^2, m_{A'}^2, M_{H_l^+}^2, M_{H_n^+}^2 \right), \quad (17)$$

$$C_2(\mu) = -\frac{1}{2} \sum_{A,A'} \sum_{l,n}^2 \frac{V_{AI}^* V_{AJ} V_{A'I}^* V_{A'J}}{16\pi^2} m_{u_A} m_{u'_A} a_R^{\dagger AI l} a_L^{AJ n} a_R^{\dagger A' I n} a_L^{A' J l} \cdot D_0 \left(m_A^2, m_{A'}^2, M_{H_l^+}^2, M_{H_n^+}^2 \right), \quad (18)$$

$$\tilde{C}_2(\mu) = -\frac{1}{2} \sum_{A,A'} \sum_{l,n}^2 \frac{V_{AI}^* V_{AJ} V_{A'I}^* V_{A'J}}{16\pi^2} m_{u_A} m_{u'_A} a_L^{\dagger AI l} a_R^{AJ n} a_L^{\dagger A' I n} a_R^{A' J l} \cdot D_0 \left(m_A^2, m_{A'}^2, M_{H_l^+}^2, M_{H_n^+}^2 \right), \quad (19)$$

$$C_4(\mu) = -\sum_{A,A'} \sum_{l,n}^2 \frac{V_{AI}^* V_{AJ} V_{A'I}^* V_{A'J}}{16\pi^2} m_{u_A} m_{u'_A} a_R^{\dagger AI l} a_L^{AJ n} a_L^{\dagger A' I n} a_R^{A' J l} \cdot D_0 \left(m_A^2, m_{A'}^2, M_{H_l^+}^2, M_{H_n^+}^2 \right), \quad (20)$$

$$C_5(\mu) = 2 \sum_{A,A'} \sum_{l,n}^2 \frac{V_{AI}^* V_{AJ} V_{A'I}^* V_{A'J}}{16\pi^2} a_L^{\dagger AI l} a_L^{AJ n} a_R^{\dagger A' I n} a_R^{A' J l} \cdot D_{00} \left(m_A^2, m_{A'}^2, M_{H_l^+}^2, M_{H_n^+}^2 \right). \quad (21)$$

These new physics contributions to the Wilson Coefficients are consistent with the ones as given in Refs. [22], the different sign and factor are due to the different definitions of the four-point integral functions. At one loop level there are no contributions to the Wilson coefficients of the operators Q_3 and $\tilde{Q}_3$.

Now, one needs to run the Wilson coefficients from the scale of new physics $\mu_t \sim M_W$ down to the low energy scale $\mu_b \sim m_b$ by using the QCD renormalization group equations. For the evolution of these coefficients, we follow Ref. [23],

$$C_r(m_b) = \sum_{i=1}^5 \sum_{s=1}^5 \left(b_i^{(r,s)} + \eta c_i^{(r,s)} \right) \eta^{a_i} C_s(\mu_t) \quad (22)$$

where we have set the new physics scale $\mu_t = m_t$ and $\eta = \alpha_s(\mu_b)/\alpha_s(m_t)$. The magic numbers a_i , $b_i^{(r,s)}$ and $c_i^{(r,s)}$ in Eq. (22) are listed in Table I.

The off-diagonal element M_{12} in the 2×2 effective Hamiltonian causes the $B^0 - \bar{B}^0$ mixing. The mass difference between the two mass eigenstates ΔM_q is described by

$$\Delta M_q = 2|M_{12}^{(q)}| \quad (23)$$

with

$$M_{12}^{(q)} = \langle \bar{B}_q^0 | \mathcal{H}_{eff}(\Delta B = 2) | B_q^0 \rangle. \quad (24)$$

In the SM, the mass splitting ΔM_q is calculated from the box diagrams of $B_q^0 - \bar{B}_q^0$ mixing, dominated by t-quark exchange. At the NLO level, one finds that [24]

$$\Delta M_q = \frac{G_F^2 M_W^2}{6\pi^2} m_{B_q} \left(\hat{B}_{B_q} f_{B_q}^2 \right) \eta_B S_0(x_t) |V_{tq} V_{tb}^*|^2 \quad (25)$$

where G_F is the Fermi constant, M_W the mass of the W boson, $S_0(x_t)$ the Inami-Lim function [21] with $x_t = m_t^2/M_W^2$. The NLO short-distance QCD correction gives $\eta_B = 0.552$, which is same for both B_d^0 and B_s^0 systems. The non-perturbative quantities $\hat{B}_{B_q}$ are the bag parameter and f_{B_q} is the B_q^0 meson decay constant [24].

In terms of the bag-parameters, the matrix elements of the operators Q_i and $\tilde{Q}_i$ are written as follows [23]:

$$\begin{aligned} \langle \bar{B}_q | \hat{Q}_1(\mu) | B_q \rangle &= \frac{1}{3} m_{B_q} f_{B_q}^2 B_1^{(q)}(\mu), \\ \langle \bar{B}_q | \hat{Q}_2(\mu) | B_q \rangle &= -\frac{5}{24} \left(\frac{m_{B_q}}{m_b(\mu) + m_q(\mu)} \right)^2 m_{B_q} f_{B_q}^2 B_2^{(q)}(\mu), \\ \langle \bar{B}_q | \hat{Q}_3(\mu) | B_q \rangle &= \frac{1}{24} \left(\frac{m_{B_q}}{m_b(\mu) + m_q(\mu)} \right)^2 m_{B_q} f_{B_q}^2 B_3^{(q)}(\mu), \\ \langle \bar{B}_q | \hat{Q}_4(\mu) | B_q \rangle &= \frac{1}{4} \left(\frac{m_{B_q}}{m_b(\mu) + m_q(\mu)} \right)^2 m_{B_q} f_{B_q}^2 B_4^{(q)}(\mu), \\ \langle \bar{B}_q | \hat{Q}_5(\mu) | B_q \rangle &= \frac{1}{12} \left(\frac{m_{B_q}}{m_b(\mu) + m_q(\mu)} \right)^2 m_{B_q} f_{B_q}^2 B_5^{(q)}(\mu), \end{aligned} \quad (26)$$

where $\hat{Q}_i(\mu)$ are the operators renormalised at the scale μ , B_i is the so-called bag factor. The matrix elements of $\tilde{Q}_{1-3}$ are the same as that of Q_{1-3} . We use the same definition of B parameters $B_i^{(q)}(\mu)$ as in Ref. [25] and find numerically that

$$\begin{aligned} B_1^{(d)}(m_b) &= 0.87(4)_{-4}^{+5}, & B_1^{(s)}(m_b) &= 0.86(2)_{-4}^{+5}, \\ B_2^{(d)}(m_b) &= 0.82(3)(4), & B_2^{(s)}(m_b) &= 0.83(2)(4), \\ B_3^{(d)}(m_b) &= 1.02(6)(9), & B_3^{(s)}(m_b) &= 1.03(4)(9), \\ B_4^{(d)}(m_b) &= 1.16(3)_{-7}^{+5}, & B_4^{(s)}(m_b) &= 1.17(2)_{-7}^{+5}, \\ B_5^{(d)}(m_b) &= 1.91(4)_{-7}^{+22}, & B_5^{(s)}(m_b) &= 1.94(3)_{-7}^{+23}. \end{aligned} \quad (27)$$

III. NUMERICAL ANALYSIS

In numerical calculations, we will use the following input parameters (all masses are in GeV)

$$\begin{aligned} m_d &= 5.4 \times 10^{-3}, & m_s &= 0.15, & m_b &= 4.6, \\ m_c &= 1.4, & \overline{m}_t(m_t) &= 165.9, & m_{B_d} &= 5.279, & m_{B_s} &= 5.367, \\ A &= 0.853, & \lambda &= 0.225, & \bar{\rho} &= 0.20 \pm 0.09, & \bar{\eta} &= 0.33 \pm 0.05, \end{aligned} \quad (28)$$

where A , λ , $\bar{\rho}$ and $\bar{\eta}$ are Wolfenstein parameters of the CKM mixing matrix.

TABLE II: The $\Delta M_q(q = d, s)$ in the SM and the T2HDM for $M_H = 300$, $\tan \beta = 10, 30, 50$ and $\delta = 0^\circ$ (a), 30° (b) and 60° (c) within 1σ range of the input hadronic parameters of JLQCD.

	SM	T2HDM		
		$\tan \beta = 10$	$\tan \beta = 30$	$\tan \beta = 50$
ΔM_d	$0.510^{+0.093}_{-0.133}$	(a) $0.511^{+0.094}_{-0.133}$	(a) $0.634^{+0.115}_{-0.162}$	(a) $1.630^{+0.289}_{-0.394}$
		(b) $0.511^{+0.095}_{-0.132}$	(b) $0.665^{+0.121}_{-0.169}$	(b) $1.737^{+0.309}_{-0.421}$
		(c) $0.510^{+0.094}_{-0.132}$	(c) $0.569^{+0.115}_{-0.145}$	(c) $1.472^{+0.260}_{-0.353}$
ΔM_s	$17.20^{+3.08}_{-2.69}$	(a) $17.15^{+3.07}_{-2.69}$	(a) $20.09^{+3.55}_{-3.11}$	(a) $44.76^{+7.61}_{-6.64}$
		(b) $17.12^{+3.06}_{-2.68}$	(b) $18.36^{+3.26}_{-2.86}$	(b) $39.32^{+6.65}_{-5.80}$
		(c) $17.07^{+3.05}_{-2.68}$	(c) $14.52^{+2.63}_{-2.30}$	(c) $26.06^{+4.27}_{-3.72}$

For the hadronic parameters $f_{B_q}^2 \hat{B}_{B_q}$, we use the values as given by the JLQCD collaboration [26],

$$\begin{aligned} f_{B_d} \hat{B}_{B_d}^{1/2}|_{\text{JLQCD}} &= (0.215^{+0.019}_{-0.030}) \text{ GeV}, \\ f_{B_s} \hat{B}_{B_s}^{1/2}|_{\text{JLQCD}} &= (0.245^{+0.021}_{-0.020}) \text{ GeV}, \end{aligned} \quad (29)$$

where the individual errors given in Ref. [26] have been added in quadrature.

In a previous paper [18], we studied the new physics contributions to the $B \rightarrow X_s \gamma$ decay, and found strong constraints on the free parameters of the considered T2HDM:

- (i) A light charged Higgs boson with a mass less than 200 GeV is excluded. For fixed $\tan \beta = 30$ and $\delta = 0^\circ$, the lower limit on M_H is $M_H \geq 300$ GeV.
- (ii) The data of $B \rightarrow X_s \gamma$ prefer a small angle δ : $\delta < 44^\circ$ for $\tan \beta = 30$ and $M_H = 400$ GeV.

Here we will consider these constraints in our choice for the free parameters of the T2HDM.

The new physics (NP) contributions to $B^0 - \bar{B}^0$ mixing are in general theoretically clean to interpret and have simple operator structure. To constrain deviations from the SM in these processes, we use the well measured physical observable ΔM_d as well as the first observation of ΔM_s as given in Eqs.(2,3) and consider the effects of the hadronic uncertainty. The theoretical predictions about the mass difference ΔM_q in the SM and T2HDM are listed in Table II for $\tan \beta = 10, 30, 50$, $M_H = 300$ GeV and $\delta = 0^\circ, 30^\circ$ and 60° . It is clear that the new physics contribution to $B^0 - \bar{B}^0$ mixing is not sensitive to the parameter δ when $\tan \beta < 20$.

In Fig. 1, we show the $\tan \beta$ dependence of ΔM_d and ΔM_s in the SM and T2HDM with the central values of hadronic parameters. The band between two horizontal dot-dashed lines shows the measured values within 3σ errors: $0.492 \leq \Delta M_d \leq 0.522$ (ps^{-1}), and $17.41 \leq \Delta M_s \leq 18.13$ (ps^{-1}). The solid horizontal lines show the central values of the SM predictions, which agree well with the data. The dominant theoretical error comes from the large uncertainty of hadronic parameter $f_{B_q} \hat{B}_{B_q}^{1/2}$. The three short-dashed curves in Fig. 1 represent the theoretical predictions in the T2HDM for $M_H = 400$ GeV and for




FIG. 1: The plots of ΔM_q vs $\tan \beta$ in the SM and T2HDM for $M_H = 400$ GeV, $\delta = 0^\circ, 30^\circ$ and 60° .

$\delta = 0^\circ, 30^\circ$ and 60° , respectively. From this figure, the upper bound on $\tan \beta$ can be read off,

$$\tan \beta \leq 25 \quad (30)$$

for $M_H = 400$ GeV and $\delta = [0^\circ, 60^\circ]$. This bound is much stronger than the one obtained from the radiative decay $B \rightarrow X_s \gamma$ [18]. After the inclusion of the effects of the uncertainties of the hadronic parameters, the upper bound on $\tan \beta$ will be changed into

$$\tan \beta \leq 30. \quad (31)$$

In Fig.2, we show the $\tan \beta$ dependence of ΔM_q in the SM and T2HDM for $\delta = 0^\circ$ and for $\tan \beta = 10, 20, 30$ and 40 , respectively. Here the central values of the input parameters are used. The same as Fig. 1, the band between two horizontal dot-dashed lines shows the measured values within 3σ errors. The solid horizontal lines also show the central values of the SM predictions. The four short-dashed curves are the theoretical predictions in the T2HDM for fixed $\delta = 0^\circ$, and for $\tan \beta = 10, 20, 30$ and 40 , respectively. It is easy to see that the data prefer a small $\tan \beta$, say $\tan \beta \leq 20$, if one assumes the existence of a light charged Higgs boson with a mass around 300 GeV. The bounds on $\tan \beta$ and M_H are indeed strongly correlated: a smaller (larger) $\tan \beta$ means a lighter (heavier) charged Higgs boson.

IV. CONCLUSIONS

In this paper, we have calculated the new physics contributions to the neutral B meson mass splitting ΔM_d and ΔM_s induced by the one-loop box diagrams involving one or two charged Higgs boson propagators in the framework of T2HDM.

By comparing the theoretical predictions with the precision data of $B_{d(s)}^0 - \bar{B}_{d(s)}^0$ mixing, strong constraints on the free parameters of T2HDM can be obtained. From the numerical results presented in last section, one can see that:




FIG. 2: The plots of ΔM_q vs M_H in the SM and T2HDM, for $\delta = 0^\circ$ and for $\tan \beta = 10, 20, 30$ and 40, respectively.

(i) For fixed $M_H = 400$ GeV and $\delta = [0^\circ, 60^\circ]$, the upper bound on $\tan \beta$ is

$$\tan \beta \leq 30 \quad (32)$$

after the inclusion of major theoretical uncertainty.

(ii) For a small $\tan \beta$, say $\tan \beta \leq 20$, a light charged Higgs boson with a mass around 300 GeV is allowed. The bounds on $\tan \beta$ and M_H are indeed strongly correlated: a smaller (larger) $\tan \beta$ means a lighter (heavier) charged Higgs boson.

Acknowledgments

One of the authors Lin-xia Lü would like to thank Professor C.-S. Huang for his valuable help. This work is partly supported by the National Natural Science Foundation of China under Grant No.10575052, and by the Specialized Research Fund for the doctoral Program of higher education (SRFDP) under Grant No. 20050319008.

-
- [1] Particle Data Group, W.M. Yao *et al.*, J. Phys. G **33** (2006) 1.
 - [2] Heavy Flavor Averaging Group, E.Barberio *et al.*, hep-ex/0603003, updated in <http://www.slac.stanford.edu/xorg/hfag>.
 - [3] CDF collaboration, A. Abulencia *et al.*, Phys. Rev. Lett. **97** (2006) 062003.
 - [4] CDF collaboration, A. Abulencia *et al.*, hep-ex/0609040.
 - [5] DØ collaboration, V. Abazov *et al.*, Phys. Rev. Lett. **97** (2006) 021802.
 - [6] UTfit Collaboration, M. Bona *et al.*, JHEP **0603** (2006) 080; CKMfitter Group, J. Charles *et al.*, Eur. Phys. J. C **41** (2005) 1 .
 - [7] L.F. Abbot, P. Sikivie, and M.B. Wise, Phys. Rev. D , **21** (1980) 1393; S.L. Glashow and E.E. Jenkins, Phys. Lett. B **196** (1987) 233; G.G. Athanasiu, P.J. Franzini, and F.J. Gilman, Phys. Rev. D **32** (1985) 3010.
 - [8] A.J. Buras, M. Jamin, and P.H. Weisz, Nucl. Phys. B **347** (1990) 491.
 - [9] A.J. Buras, S. Jäger, and J. Urban, Nucl. Phys. B **605** (2001) 600.
 - [10] J. Urban, F. Krauss, U. Jentschura, and G. Soff, Nucl. Phys. B **523** (1998) 40.
 - [11] C.S. Huang and J.T. Li, Int.J.Mod.Phys.A **20** (2005)161
 - [12] A. Grant, Phys. Rev. D **51** (1995) 207; D. Atwood, L. Reina and A. Soni, Phys. Rev. D **55** (1997) 3156; D. Bowser-Chao, K. Cheung, and W.-Y. Keung, Phys. Rev. D **59** (1999) 115006.
 - [13] Z.J. Xiao and L.B. Guo, Phys. Rev. D **69** (2004) 014002.
 - [14] G. Isidori and P. Paradisi, Phys. Lett. B **639** (2006) 499; K. Cheung, C.-W. Chiang, N.G. Deshpande, and J. Jiang, hep-ph/0604223; S. Baek, hep-ph/0605182; S. Baek, J.H. Jeon, and C.S. kim, Phys. Lett. B **641** (2006) 183; B. Dutta, and Y. Mimura, hep-ph/0607147; S. Chang, C.S. Kim, and J. Song, hep-ph/0607313; F.J. Botella, G.C. Branco, and M. Nebot, hep-ph/0608100; J.K. Parry, hep-ph/0608192; S. Nandi, and J.P. Saha, hep-ph/0608341; Gao Xiang-dong, C.S. Li, L.L. Yang, hep-ph/0609269.
 - [15] A. Das and C. Kao, Phys. Lett. B **372** (1996) 106.
 - [16] K. Kiers, A. Soni and G.H. Wu, Phys. Rev. D **59** (1999) 096001; G.H. Wu and A. Soni, Phys. Rev. D **62** (2000) 056005.
 - [17] K. Kiers, A. Soni and G-H. Wu, Phys. Rev. D , **62** (2000) 116004.
 - [18] Zhen-jun Xiao and Lin-xia Lü, Phys. Rev. D **74** (2006) 034016.
 - [19] W.-S. Hou, Phys. Lett. B **296** (1992) 179 ; M. Luke and M.J. Savage, Phys. Lett. B **307** (1993) 387.
 - [20] J. Rosiek, Phys. Rev. D **41** (1990) 3464, Erratum hep-ph/9511250 (unpublished).
 - [21] T. Inami and C. Lim, Prog.Theor.Phys. **65** (1981) 297.
 - [22] A.J. Buras, P.H. Chankowski, J. Rosiek, and L. Slawianowska, Nucl. Phys. B **619** (2001) 434.
 - [23] D. Bećirević *et al.*, Nucl. Phys. B **634** (2002) 105.
 - [24] G. Buchalla, A.J. Buras, and M.E. Lautenbacher, Rev. Mod. Phys. **68**, 1125 (1996).
 - [25] D. Bećirević *et al.*, Nucl. Phys. Proc. Suppl. **106** (2002) 385; JHEP **0204** (2002) 025.
 - [26] S. Aoki *et al.*, JLQCD Collaboration, Phys. Rev. Lett. **91** (2003) 212001.